VOL. 471 JOURNAL OF CHROMATOGRAPHY JUNE 2, 1989

CONTENTS

HONOUR VOLUME ON THE OCCASION OF THE 65TH BIRTHDAY OF R. P. W. SCOTT

Historical

Profile of R. P. W. Scott
by D. E. Martire 1

Critical reviews

Ribonucleoside analysis by reversed-phase high-performance liquid chromatography
by C. W. Gehrke and K. C. Kuo (Columbia, MO, U.S.A.) 3

Gas–solid chromatography on open-tubular columns: an isotope effect
by M. Mohnke and J. Heybey (Leipzig, G.D.R.) 37

Basic equations in continuous gas extraction and their application to headspace analysis
by A. G. Vitenberg and B. V. Ioffe (Leningrad, U.S.S.R.) 55

Theory

Theory of open-tube chromatography: and exact proof of Golay's equations
by A. A. Clifford (Leeds, U.K.) 61

Application of a unified molecular theory to gas–liquid chromatography
by D. E. Martire (Washington, DC, U.S.A.) 71

Comparison between the conditions for solute focussing by the static and dynamic solvent effects under ideal conditions
by P. J. Apps and V. Pretorius (Pretoria, South Africa) 81

Gas chromatography

Comparison of solvent models for characterizing stationary phase selectivity in gas chromatography
by S. K. Poole, B. R. Kersten and C. F. Poole (Detroit, MI, U.S.A.) 91

Determination of the gas chromatographic performance characteristics of several graphitized carbon blacks
by W. R. Betz and W. R. Supina (Bellefonte, PA, U.S.A.) 105

Characteristics of glass based packings as a support in chromatography
by L. S. Green and W. Bertsch (Tuscaloosa, AL, U.S.A.) 113

New method for measuring obstructive factors and porosity using gas chromatographic instrumentation
by N. A. Katsanos and Ch. Vassilakos (Patras, Greece) 123

Use of gas chromatography for a study of crown ethers
by A. Kohoutová, E. Smolková-Keulemansová and L. Feltl (Prague, Czechoslovakia) . . 139

Isomerization of propyne to propadiene. Studies by gas chromatography
by P. M. Lyne and C. S. G. Phillips (Oxford, U.K.) 145

Sample preparation by means of a supported liquid membrane for the determination of chlorophenoxyalkanoic acids
by G. Nilvé, G. Audunsson and J. Å. Jönsson (Lund, Sweden) 151

Quantitative high-resolution gas chromatography and mass spectrometry of toxaphene residues in fish samples
by F. I. Onuska and K. A. Terry (Burlington, Canada) 161

Determination of sensorial active trace compounds by multi-dimensional gas chromatography combined with different enrichment techniques
by S. Nitz, H. Kollmannsberger and F. Drawert (Freising-Weihenstephan, F.R.G.) . . . 173

Relative gas–liquid chromatographic retention factors of trimethylsilyl ethers of diradylglycerols on polar capillary columns
by J. J. Myher and A. Kuksis (Toronto, Canada) 187

Gas chromatographic retention of carbohydrate trimethylsilyl ethers. III. Ketohexoses
by A. Garciá-Raso (Palma de Mallorca, Spain), and M. Fernández-Díaz, M. I. Páez, J. Sanz and I. Martínez-Castro (Madrid, Spain) 205

Dispersion and selectivity indices in gas chromatography. Part II. Studies of homologous carbonyl and carboxyl compounds
by M. B. Evans (Hatfield, U.K.) and J. K. Haken (Kensington, Australia) 217

Quantitative relationships between the structure of alkylbenzenes and their gas chromatographic retention on stationary phases with different polarity
by N. Dimov (Sofia, Bulgaria) and Ov. Mekenyan (Burgas, Bulgaria) 227

Effect of repeated cross-linking of SE-54 stationary phase film on the chromatographic properties of capillary columns
by K. Janák, M. Horká and K. Tesařík (Brno, Czechoslovakia) 237

Non-linearity of the plot of log (adjusted retention time) *versus* carbon number of *n*-alkanes in series-coupled gas chromatographic columns
by T. Maurer, Th. Welsch and W. Engewald (Leipzig, G.D.R.) 245

Interface adsorption and reproducibility of retention indices in glass capillary columns with dimethylpolysiloxane stationary phases cross-linked by γ-irradiation
by P. Farkaš, L. Soják and M. Kováč (Bratislava, Czechoslovakia) and J. Janák (Brno, Czechoslovakia) 251

Liquid chromatography

Determination of the enantiomers of α-H-α-amino acids, α-alkyl-α-amino acids and the corresponding acid amides by high-performance liquid chromatography
by A. Duchateau, M. Crombach, J. Kamphuis, W. H. J. Boesten, H. E. Schoemaker and E. M. Meijer (Geleen, The Netherlands) 263

Improved chiral stationary phase for the separation of the enantiomers of chiral acids as their anilide derivatives
by W. H. Pirkle and J. E. McCune (Urbana, IL, U.S.A.) 271

Chromatographic evaluation of oligomeric C_8 reversed phases for the use in high-performance liquid chromatography
by S. O. Akapo, A. Furst, T. M. Khong and C. F. Simpson (London, U.K.) 283

Mass detection limits achieved with a commercially available fluorimeter in micro high-performance liquid chromatography
by T. Takeuchi, T. Asano and D. Ishii (Nagoya, Japan) 297

Simultaneous photometric and conductivity detection for microcolumn liquid chromatography
by M. Janeček and K. Šlais (Brno, Czechoslovakia) 303

Effect of the position of the laser beam focal point on a capillary flow-through cell on the signal-to-noise ratio for a fluorimetric detector in capillary column liquid chromatography
by T. Tsuda (Nagoya, Japan) and H. Noda (Kyoto, Japan) 311

Trace enrichment of pyrimidine nucleobases, 5-fluorouracil and bromacil on a silver-loaded thiol stationary phase with on-line reversed-phase high-performance liquid chromatography
by C. Lipschitz, H. Irth, G. J. de Jong, U. A. Th. Brinkman and R. W. Frei (Amsterdam, The Netherlands) 321

Quantitative microscale liquid chromatography of piperine in pepper and pepper extracts
by M. Verzele, F. Van Damme and G. Schuddink (Ghent, Belgium) and P. Vyncke (Eke, Belgium) 335

Supercritical fluid chromatography

Effect of pressure drop across the column on average densities and capacity factors in supercritical fluid chromatography
by K. D. Bartle, T. Boddington, A. A. Clifford and G. F. Shilstone (Leeds, U.K.) 347

Adsorption isotherms on silica for methanol and 1-hexanol modifiers from supercritical carbon dioxide
by C. H. Lochmüller and L. P. Mink (Durham, NC, U.S.A.) 357

Other chromatographic techniques and combinations

Aqueous size-exclusion chromatography of anionic and non-ionic water-soluble polymers on silica gel with bonded hydrophilic groups
by S. Mori (Mie, Japan) 367

Overpressured multi-layer chromatography
by E. Tyihák, E. Mincsovics and T. J. Székely (Budapest, Hungary) 375

Laser microprobe mass spectrometry of selected compounds directly from normal phase high-performance thin-layer plates
by R. W. Finney and H. Read (Sunbury-on-Thames, U.K.) 389

Determination of free and esterified sterols and of wax esters in oils and fats by coupled liquid chromatography–gas chromatography
by K. Grob and M. Lanfranchi (Zürich, Switzerland) and C. Mariani (Milan, Italy) . . . 397

Retention behaviour of β-carbolines in normal-phase chromatography. Silica and amino phases in high-performance and thin-layer chromatography
by M. C. Pietrogrande, F. Dondi, P. A. Borea and C. Bighi (Ferrara, Italy) 407

Capillary electrophoresis

Zone broadening due to sample injection in capillary zone electrophoresis
by E. Grushka and R. M. McCormick (Wilmington, DE, U.S.A.) 421

Capillary zone electrophoretic separations of proteins in polyethylene glycol-modified capillaries
by G. J. M. Bruin, J. P. Chang, R. H. Kuhlman, K. Zegers, J. C. Kraak and H. Poppe (Amsterdam, The Netherlands) 429

Research notes

Picolinic acid: a mobile phase additive for improved chromatography of metal-chelating heterocyclic acids and β-diketones
by D. W. Roberts, R. J. Ruane and I. D. Wilson (Macclesfield, U. K.) 437

Trace/ultratrace analyses of unstable compounds. Investigations on hydrazobenzene and azobenzene
by S. Ahuja, G. Thompson and J. Smith (Suffern, NY, U.S.A.) 443

Letter to the Editor

On some nomenclature in chromatography
by O. K. Guha (Bihar, India) 447

Author Index 449